

Inequalities involving Central Cevians

STANLEY RABINOWITZ

545 Elm St Unit 1, Milford, New Hampshire 03055, USA

e-mail: stan.rabinowitz@comcast.net

web: <http://www.StanleyRabinowitz.com/>

Abstract. A cevian of a triangle is a line segment that extends from a vertex of the triangle to a point on the opposite side. A cevian that passes through a triangle center is called a central cevian. There are a number of inequalities known concerning central cevians. For example, if m_a , m_b , and m_c are the lengths of the medians of a triangle, then it is known that

$$27r^2 \leq m_a^2 + m_b^2 + m_c^2 \leq \frac{27}{4}R^2$$

where r is the inradius of the triangle and R is its circumradius. We use a computer to discover and prove similar inequalities for other central cevians. For example, if f_a , f_b , and f_c are the lengths of the Feuerbach cevians of a triangle, then

$$\frac{7}{8}s^2 \leq f_a^2 + f_b^2 + f_c^2 \leq \frac{64}{7}R^2$$

where s is the semiperimeter of the triangle.

Keywords. triangle centers, inequalities, computer-discovered mathematics, cevians, Mathematica.

Mathematics Subject Classification (2020). 51M04, 51-08.

1. INTRODUCTION

There are many notable points associated with a triangle, such as the incenter, centroid, circumcenter, and orthocenter, These are special cases of *triangle centers* as defined by Clark Kimberling in [3]. A *cevian* of a triangle is a line segment that extends from a vertex of the triangle to a point on the opposite side. A cevian that passes through a triangle center is called a *central cevian*. The cevian from vertex A is called the *A-cevian*. The other cevians are named similarly.

Let X_n denote the n th named triangle center as cataloged in the Encyclopedia of Triangle Centers [4]. Let $|PQ|$ denote the length of the line segment PQ .

The cevians through X_n will be named AA_n , BB_n , and CC_n as shown in Figure 1.

¹This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.




FIGURE 1. Cevians through X_n

2. THE DATA

We use barycentric coordinates in this study. The barycentric coordinates for triangle centers X_1 through X_{12} in terms of the sides of the triangle, a , b , and c , are shown in Table 1. Only the first barycentric coordinate is given, because if $f(a, b, c)$ is the first barycentric coordinate for a point P , then the barycentric coordinates for P are

$$\left(f(a, b, c) : f(b, c, a) : f(c, a, b) \right).$$

These were derived from [4].

TABLE 1. Barycentric coordinates for the first 12 centers

n	First barycentric coordinate for X_n
1	a
2	1
3	$a^2(a^2 - b^2 - c^2)$
4	$(a^2 + b^2 - c^2)(a^2 - b^2 + c^2)$
5	$c^4 - a^2b^2 + b^4 - a^2c^2 - 2b^2c^2$
6	a^2
7	$(a + b - c)(a - b + c)$
8	$a - b - c$
9	$a(a - b - c)$
10	$b + c$
11	$(b - c)^2(-a + b + c)$
12	$(a + b - c)(a - b + c)(b + c)^2$

We will find inequalities that involve the squares of the lengths of central cevians and other elements of a triangle, as listed in Table 2.

TABLE 2. Elements of a triangle

symbol	Description
a, b, c	the sides of the triangle
K	the area of the triangle
r	the inradius of the triangle
R	the circumradius of the triangle
s	the semiperimeter of the triangle

To find the distance between two points, we used the following formula which comes from [2].

Proposition 1 (Distance Formula). *Given two points $P = (u_1, v_1, w_1)$ and $Q = (u_2, v_2, w_2)$ in normalized barycentric coordinates. Denote $x = u_1 - u_2$, $y = v_1 - v_2$ and $z = w_1 - w_2$. Then the distance between P and Q is*

$$\sqrt{-a^2yz - b^2xz - c^2xy}.$$

To find the length of a cevian of a triangle, we proceed as follows. Set up a barycentric coordinate system with $\triangle ABC$ as the reference triangle, so that $A = (1 : 0 : 0)$, $B = (0 : 1 : 0)$, and $C = (0 : 0 : 1)$. Let P be an arbitrary point in the plane other than A . Let the barycentric coordinates for P be $(p : q : r)$. Let AP meet BC at A' (Figure 2). Then it is straightforward to show that the barycentric coordinates for A' are $(0 : q : r)$.




FIGURE 2. Barycentric Coordinates

Using Proposition 1, we get the following result.

Proposition 2 (Cevian Length). *Let P be a point in the plane of $\triangle ABC$ with trilinear coordinates $(p : q : r)$. Let AP meet BC at A' . Then*

$$|AA'| = \frac{\sqrt{b^2r(q+r) + c^2q(q+r) - a^2qr}}{q+r}.$$

Using Proposition 2 and Table 1, we can find the length of the A -cevian that passes through the point X_n . Table 3 shows the lengths for n ranging from 1 to 12, where $K = \sqrt{s(s-a)(s-b)(s-c)}$ and $s = (a+b+c)/2$.

TABLE 3. Cevian lengths for the first 12 centers

n	Square of length of A -cevia passing through X_n
1	$bc \left(1 - \frac{a^2}{(b+c)^2} \right)$
2	$\frac{1}{4} (2(b^2 + c^2) - a^2)$
3	$-\frac{a^2 b^2 c^2 (a^4 - 2a^2(b^2 + c^2) + (b^2 - c^2)^2)}{((b^2 - c^2)^2 - a^2(b^2 + c^2))^2}$
4	$\frac{4K^2}{a^2}$
5	$\frac{16K^2 (a^6 - 3a^4 b^2 - 3a^4 c^2 + 3a^2 b^4 + 3a^2 b^2 c^2 + 3a^2 c^4 - b^6 + b^4 c^2 + b^2 c^4 - c^6)}{(2a^4 - 3a^2 b^2 - 3a^2 c^2 + b^4 - 2b^2 c^2 + c^4)^2}$
6	$\frac{b^2 c^2 (2(b^2 + c^2) - a^2)}{(b^2 + c^2)^2}$
7	$-\frac{a^3 + a(-3b^2 + 2bc - 3c^2) + 2(b-c)^2(b+c)}{4a}$
8	$\frac{-a^3 + a(3b^2 - 2bc + 3c^2) + 2(b-c)^2(b+c)}{4a}$
9	$-\frac{bc(a^4 - 2a^2(b^2 + c^2) + (b-c)^4)}{((b-c)^2 - a(b+c))^2}$
10	$\frac{a^2(-(a+b))(a+c) + b^2(a+b)(2a+b+c) + c^2(a+c)(2a+b+c)}{(2a+b+c)^2}$
11	$\frac{(b^2 x + c^2 y)(x+y) - a^2 xy}{(x+y)^2}$ where $x = (a-b)^2(a+b-c)$ $y = (a-c)^2(a-b+c)$
12	$\frac{xy(-a^2 + b^2 + c^2) + b^2 y^2 + c^2 x^2}{(x+y)^2}$ where $x = (a+c)^2(a+b-c)$ $y = (a+b)^2(a-b+c)$

3. MAIN RESULTS

Notation. The symbol S_n represents the sum of the squares of the lengths of the cevians of $\triangle ABC$ that pass through triangle center X_n . In other words,

$$S_n = |AA_n|^2 + |BB_n|^2 + |CC_n|^2.$$

For example, if $n = 2$, then the cevians are medians and $S_2 = m_a^2 + m_b^2 + m_c^2$.

Conventions. In this section, all inequalities listed are best possible.

The inequality $S_n \leq k_0 f(a, b, c)$ is said to be *best possible* if there is no constant k with $k < k_0$ such that $S_n \leq k f(a, b, c)$ is true for all triangles.

The inequality $k_0 f(a, b, c) \leq S_n$ is said to be *best possible* if there is no constant k with $k > k_0$ such that $k f(a, b, c) \leq S_n$ is true for all triangles.

If no upper bound is listed for S_n with respect to $f(a, b, c)$, this means that there is no constant k such that $S_n \leq k f(a, b, c)$ is true for all triangles.

If no lower bound is listed for S_n with respect to $f(a, b, c)$, this means that there is no constant $k > 0$ such that $k f(a, b, c) \leq S_n$ is true for all triangles.

Methodology. The best constants for all inequalities were found using Mathematica and Algorithm K from [8]. Since all computations were performed using exact symbolic algebra (as opposed to numerical approximations), these computer calculations constitute proofs that the inequalities are correct.

Theorem 1. *The following inequalities are true for all triangles.*

$$\begin{aligned} 27r^2 &\leq S_1 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_2 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_3 \\ 27r^2 &\leq S_4 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_5 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_6 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_7 \leq \frac{27}{4}R^2 \\ 27r^2 &\leq S_8 < 12R^2 \\ 27r^2 &\leq S_9 < \frac{68}{9}R^2 \\ 27r^2 &\leq S_{10} < \frac{68}{9}R^2 \\ k_1r^2 &\leq S_{11} \leq \frac{64}{7}R^2 \\ 27r^2 &\leq S_{12} \leq \frac{27}{4}R^2 \end{aligned}$$

where $k_1 \approx 30.91612615$ is the positive root of $x^3 - 32x^2 + 48x - 448$.

Equality occurs when the triangle is equilateral, except in the following cases.

For $27r^2 \leq S_3$, $S_5 \leq \frac{27}{4}R^2$, and $27r^2 \leq S_5$, equality occurs when the sides of the triangle are proportional to 1, 1, and $\sqrt{3}$.

For $S_{11} \leq \frac{64}{7}R^2$, equality occurs when the sides of the triangle are proportional to 1, 1, and $2\sqrt{\frac{3}{7}}$.

For $k_1r^2 \leq S_{11}$, equality occurs when the sides of the triangle are proportional to 1, 1, and the positive root of $7x^3 + 2x^2 + 4x - 8$.

Lemma 1. Let A' be a point in the interior of side BC of $\triangle ABC$. Let $|AB| = c$, $|AC| = b$ and $|AA'| = x_a$ (Figure 3). Then

$$h_a \leq x_a < \max(b, c)$$

where h_a is the length of the altitude from A .




FIGURE 3. Cevian from A

Proof. Let H be the foot of the altitude from A (Figure 4). By the Pythagorean Theorem, it can be seen that the closer A' gets to H , the smaller x_a gets. The minimum value of x_a is h_a and the maximum value for x_a is the larger of b and c . \square




FIGURE 4. Cevian from A

Proposition 3. Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then

$$27r^2 \leq x < 12R^2.$$

Proof. We will prove a more general result. Let x_a be the length of any interior cevian from vertex A of $\triangle ABC$. (An *interior cevian* meets the opposite side at an interior point of that side.) Define x_b and x_c similarly. Note that the three cevians need not all pass through the same point P . Then we will show that

$$(1) \quad 27r^2 \leq x_a^2 + x_b^2 + x_c^2 < 12R^2.$$

By Lemma 1, $x_a \geq h_a$. Similarly for x_b and x_c . Thus,

$$x_a^2 + x_b^2 + x_c^2 \geq h_a^2 + h_b^2 + h_c^2.$$

But

$$h_a^2 + h_b^2 + h_c^2 \geq 27r^2$$

from inequality $27r^2 \leq S_4$ of Theorem 1. This proves the left side of Equation (1). Without loss of generality, we can assume that $a \leq b \leq c$. By Lemma 1, we have $x_a < c$, $x_b < c$, and $x_c < b$. Thus

$$(2) \quad x_a^2 + x_b^2 + x_c^2 < b^2 + 2c^2.$$

The right side of Equation (1) will then be true if we can prove that $b^2 + 2c^2 < 12R^2$. This inequality is not homogeneous, so we cannot use the methods of [8]. Instead, we use the `Simplify` command in Mathematica. The formula for R in terms of a , b , and c is well known, namely

$$R = \frac{abc}{4K}$$

where K is the area of $\triangle ABC$. We thus issue the following Mathematica commands.

```
s = (a+b+c)/2;
K = Sqrt[s(s-a)(s-b)(s-c)];
R = a*b*c/(4K);
inequality = b^2+2c^2 < 12R^2;
triangCondition = a>0 && b>0 && c>0 && a+b>c && b+c>a && c+a>b;
Simplify[inequality, triangCondition]
```

Mathematica responds with `True`, indicating that the inequality is correct. Note that we did not need the condition $a \leq b \leq c$. This concludes the proof of the right side of Equation (1). \square

The constants in Proposition 3 are best possible as can be seen by the inequality for S_8 in Theorem 1.

Continuing with Algorithm K, we get the following results.

Theorem 2. *The following inequalities are true for all triangles.*

$$\begin{aligned} \frac{8}{9}s^2 &< S_1 \leq s^2 \\ s^2 &\leq S_2 < \frac{3}{2}s^2 \\ S_4 &\leq s^2 \\ S_5 &\leq \frac{33}{25}s^2 \\ \frac{18}{25}s^2 &< S_6 \leq s^2 \\ \frac{1}{2}s^2 &< S_7 \leq s^2 \\ s^2 &\leq S_8 < 3s^2 \\ s^2 &\leq S_9 < 2s^2 \\ s^2 &\leq S_{10} < \frac{17}{9}s^2 \\ \frac{7}{8}s^2 &\leq S_{11} \leq 2s^2 \\ \frac{1}{2}s^2 &< S_{12} \leq s^2 \end{aligned}$$

Equality occurs when the triangle is equilateral, except in the following cases.

For $S_5 \leq \frac{33}{25}s^2$, equality occurs when the sides of the triangle are proportional to 1, 1, and $\sqrt{3}$.

For $\frac{7}{8}s^2 \leq S_{11}$, equality occurs when the sides of the triangle are proportional to 1, 1, and $\frac{2}{7}$.

Proposition 4. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$x < 3s^2.$$

Proof. This inequality follows from Equation (2) and the fact that the Mathematica code

```
inequality = b^2+2c^2 < 3s^2;
Simplify[inequality, triangCondition]
```

returns `True`. □

The constant “3” is best possible as can be seen from the inequality for S_8 in Theorem 2.

Lemma 2. *Let A' be a point in the interior of side BC of $\triangle ABC$. Let $|AB| = c$, $|AC| = b$ and $|AA'| = x_a$ (Figure 3). Then*

$$x_a \geq \min(b, c) \cos \frac{A}{2}.$$

Proof. In [5], it is shown that

$$x_a \geq (kb + k'c) \cos \frac{A}{2}$$

where $k = |BA'|/|A'C|$ and $k' = 1 - k$. The function $f(k) = kb + (1 - k)c$ is a linear function of k over the interval $[0, 1]$. It takes on all values from $\min(b, c)$ to $\max(b, c)$. Therefore, when $b > 0$, $c > 0$, and $0 \leq k \leq 1$, we must have

$$kb + (1 - k)c \geq \min(b, c).$$

The result now follows. □

Proposition 5. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$\frac{1}{2}s^2 < x.$$

Proof. By Lemma 2, we have

$$x_a^2 \geq \min(b^2, c^2) \cos^2 \frac{A}{2}.$$

From the half-angle formula for cosine,

$$\cos^2 \frac{A}{2} = \frac{1 + \cos A}{2},$$

and from the Law of Cosines,

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc},$$

we see that

$$x_a^2 \geq \min(b^2, c^2) \frac{(b + c - a)(b + c + a)}{4bc}$$

with similar formulas for x_b and x_c .

Without loss of generality, assume that $a \leq b \leq c$. Then

$$(3) \quad x = x_a^2 + x_b^2 + x_c^2 \geq b^2 \frac{(b + c - a)s}{2bc} + a^2 \frac{(c + a - b)s}{2ca} + a^2 \frac{(a + b - c)s}{2ab}.$$

Using this value for x in terms of a , b , and c , and the definitions of s and `triangCondition` from the proof of Proposition 3, we issue the following Mathematica commands.

```
inequality = x > (1/2)s^2;
Simplify[inequality, triangCondition]
```

The response of `True` proves the inequality. □

The proof shows that the result is true for any three internal cevians. They do not necessarily have to all pass through the same point P .

The constant “1/2” is best possible as can be seen from the inequality for S_7 in Theorem 2.

Related results can be found in Theorem 14.11 of [1, p. 124] and Theorem 7.22 of [6, p. 337].

Continuing with Algorithm K, we get the following results.

Theorem 3. *The following inequalities are true for all triangles.*

$$\begin{aligned} \frac{27}{2}rR &\leq S_1 \leq \frac{3}{2}\sqrt{3}Rs \\ \frac{27}{2}rR &\leq S_2 < 3Rs \\ S_4 &\leq \frac{3}{2}\sqrt{3}Rs \\ S_5 &\leq \frac{66}{25}Rs \\ \frac{288}{25}rR &< S_6 \leq \frac{3}{2}\sqrt{3}Rs \\ 8rR &< S_7 \leq \frac{3}{2}\sqrt{3}Rs \\ \frac{27}{2}rR &\leq S_8 < 6Rs \\ \frac{27}{2}rR &\leq S_9 < \frac{34}{9}Rs \\ \frac{27}{2}rR &\leq S_{10} < \frac{34}{9}Rs \\ k_2rR &\leq S_{11} \leq k_3Rs \\ 8rR &< S_{12} \leq \frac{3}{2}\sqrt{3}Rs \end{aligned}$$

where $k_2 \approx 14.12657721$ is the positive root of $2x^3 - 5x^2 - 256x - 1024$ and $k_3 \approx 3.737553924$ is the largest positive root of $4x^6 + 20117x^4 - 356864x^2 + 1048576$. Equality occurs when the triangle is equilateral, except for $S_5 \leq \frac{66}{25}Rs$, where equality occurs when the sides of the triangle are proportional to 1, 1, and $\sqrt{3}$.

Proposition 6. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$8rR < x < 6Rs.$$

Proof. The right side of the inequality follows from Equation (2) and the fact that the Mathematica code

```
inequality = b^2+2c^2 < 6R*s;
Simplify[inequality, triangCondition]
```

returns `True`.

The left side of the inequality follows from the fact that the Mathematica code

```
r = K/s;
inequality = x > 8rR;
Simplify[inequality, triangCondition]
```

returns `True`, where x is given by Equation (3). □

The constants in Proposition 6 are best possible as can be seen from the inequalities for S_7 and S_8 in Theorem 3.

Continuing with Algorithm K, we get the following results.

Theorem 4. *The following inequalities are true for all triangles.*

$$\begin{aligned} 3\sqrt{3}K &\leq S_1 \\ 3\sqrt{3}K &\leq S_2 \\ 3\sqrt{3}K &\leq S_6 \\ 3\sqrt{3}K &\leq S_7 \\ 3\sqrt{3}K &\leq S_8 \\ 3\sqrt{3}K &\leq S_9 \\ 3\sqrt{3}K &\leq S_{10} \\ 4\sqrt{2}K &\leq S_{11} \\ 3\sqrt{3}K &\leq S_{12} \end{aligned}$$

Equality occurs when the triangle is equilateral, except for $4\sqrt{2}K \leq S_{11}$, where equality occurs when the sides of the triangle are proportional to 1, 1, and $\frac{2}{3}$.

Proposition 7. *Let x_a be the length of an internal A -cevia in $\triangle ABC$. Define x_b and x_c similarly. (The three cevians need not concur.) Let $x = x_a^2 + x_b^2 + x_c^2$. Then*

$$x \geq k_7 K$$

where $k_7 \approx 4.319536403$ is the positive real root of $x^{26} + 279x^{24} + 26353x^{22} + 1287331x^{20} + 29550479x^{18} - 84430591x^{16} - 19873132241x^{14} - 177553607339x^{12} + 3995469783904x^{10} - 20956237447808x^8 + 24820097419264x^6 - 17358828744704x^4 + 5114979942400x^2 - 1274019840000$ and is the best possible constant.

Proof. The following Mathematica code proves this result.

```
s = (a+b+c)/2;
K = Sqrt[s(s-a)(s-b)(s-c)];
expression = x/K;
Minimize[expression, triangCondition, {a,b,c}]
```

where x is given by Equation (3). □

If the 3 cevians concur, then Theorem 4 would suggest that $x_a^2 + x_b^2 + x_c^2 \geq 3\sqrt{3}K$. However, this is not the case. Figure 5 shows an example where

$$\frac{x_a^2 + x_b^2 + x_c^2}{K} \approx 4.95030 < 3\sqrt{3}.$$




FIGURE 5. Three concurrent cevians with $(x_a^2 + x_b^2 + x_c^2) < 3\sqrt{3}K$.

In this figure, found using Geometer's Sketchpad, $a = 3$, $b = 105/32$, $c = 39/8$, $|B'C| \approx 2.10706$, $|AC'| \approx 2.76794$, $|BA'| \approx 2.93549$, $|CA'| \approx 0.06451$, $|AB'| \approx 3.22726$, $|CB'| \approx 0.05399$, $x_a \approx 3.29496$, $x_b \approx 3.01143$, $x_c \approx 1.98276$, and $(x_a^2 + x_b^2 + x_c^2)/K \approx 4.95030$.

Noting that $4.95030^2 \approx 24.5$ suggests the following conjecture.

Conjecture 1. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$x \geq \frac{7}{2}\sqrt{2}K.$$

4. RELATED RESULTS

In this section, the inequalities are all best possible, however, we omit inequalities involving S_n for $n = 3, 5, 11,$ or 12 . The compute power available to us was insufficient to find many of the inequalities involving these central cevians. The results were found using Algorithm K.

Theorem 5. *The following inequalities are true for all triangles.*

$$\begin{aligned} \frac{1}{2}(a^2 + b^2 + c^2) < S_1 &\leq \frac{3}{4}(a^2 + b^2 + c^2) \\ S_2 &= \frac{3}{4}(a^2 + b^2 + c^2) \\ S_4 &\leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{12}{25}(a^2 + b^2 + c^2) < S_6 &\leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{1}{3}(a^2 + b^2 + c^2) < S_7 &\leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_8 < \frac{3}{2}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_9 < \frac{4}{3}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_{10} < \frac{17}{18}(a^2 + b^2 + c^2) \end{aligned}$$

Equality occurs when the triangle is equilateral.

Proposition 8. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$\frac{1}{3}(a^2 + b^2 + c^2) < x < \frac{1}{3}(3 + \sqrt{3})(a^2 + b^2 + c^2).$$

The proof uses Mathematica in the same way as in the proof of Proposition 6 and is omitted.

The constant “1/3” on the left side of the inequality in Proposition 8 is best possible as can be seen from the inequality for S_7 in Theorem 5. To show that the constant “ $\frac{1}{3}(3 + \sqrt{3})$ ” on the right side is best possible, we recall Equation (2), and then issue the following Mathematica command.

```
expression = (b^2+2c^2)/(a^2+b^2+c^2);
Maximize[expression, triangCondition, {a,b,c}]
```

Mathematica returns the maximum $\frac{1}{3}(3 + \sqrt{3})$ and states that the maximum occurs for the degenerate triangle with sides $1, 1 + \sqrt{3},$ and $2 + \sqrt{3}$.

Continuing with Algorithm K, we get the following results.

Theorem 6. *The following inequalities are true for all triangles.*

$$\begin{aligned} \frac{32}{45}(ab + bc + ca) &< S_1 < ab + bc + ca \\ \frac{3}{4}(ab + bc + ca) &\leq S_2 < \frac{3}{2}(ab + bc + ca) \\ S_4 &< ab + bc + ca \\ \frac{72}{125}(ab + bc + ca) &< S_6 < ab + bc + ca \\ \frac{2}{5}(ab + bc + ca) &< S_7 < ab + bc + ca \\ \frac{3}{4}(ab + bc + ca) &\leq S_8 < 3(ab + bc + ca) \\ \frac{3}{4}(ab + bc + ca) &\leq S_9 < \frac{17}{9}(ab + bc + ca) \\ \frac{3}{4}(ab + bc + ca) &\leq S_{10} < \frac{17}{9}(ab + bc + ca) \end{aligned}$$

Equality occurs when the triangle is equilateral.

Proposition 9. *Let P be a point inside $\triangle ABC$. Let x be the sum of the squares of the lengths of the cevians through P . Then*

$$\frac{2}{5}(ab + bc + ca) < x < 3(ab + bc + ca).$$

The proof uses Mathematica in the same way as in the proof of Proposition 6 and is omitted.

The constants in the inequality in Proposition 9 are best possible as can be seen from the inequalities for S_7 and S_8 in Theorem 6.

Continuing with Algorithm K, we get the following results.

Theorem 7. *The following inequalities are true for all triangles.*

$$\begin{array}{llll}
 S_1 \leq S_2 & & & S_9 \leq \frac{9}{4}S_1 \\
 S_1 \leq \frac{100}{81}S_6 & S_4 \leq S_1 & S_7 \leq S_1 & S_9 \leq \frac{16}{9}S_2 \\
 S_1 \leq \frac{16}{9}S_7 & S_4 \leq S_2 & S_7 \leq S_2 & S_9 \leq \frac{25}{9}S_6 \\
 S_1 \leq S_8 & S_4 \leq S_6 & S_7 \leq k_4S_6 & S_9 \leq 4S_7 \\
 S_1 \leq S_9 & S_4 \leq S_7 & S_7 \leq S_8 & S_9 \leq S_8 \\
 S_1 \leq S_{10} & S_4 \leq S_8 & S_7 \leq S_9 & S_9 \leq \frac{25}{16}S_{10} \\
 & S_4 \leq S_9 & S_7 \leq S_{10} & \\
 & S_4 \leq S_{10} & & \\
 \\
 S_2 \leq \frac{3}{2}S_1 & & S_8 \leq 3S_1 & S_{10} \leq \frac{17}{9}S_1 \\
 S_2 \leq \frac{25}{16}S_6 & S_6 \leq S_1 & S_8 \leq 2S_2 & S_{10} \leq \frac{34}{27}S_2 \\
 S_2 \leq \frac{9}{4}S_7 & S_6 \leq S_2 & S_8 \leq 3S_6 & S_{10} \leq \frac{17}{9}S_6 \\
 S_2 \leq S_8 & S_6 \leq \frac{36}{25}S_7 & S_8 \leq 4S_7 & S_{10} \leq \frac{64}{25}S_7 \\
 S_2 \leq S_9 & S_6 \leq S_8 & S_8 \leq \frac{27}{17}S_9 & S_{10} \leq S_8 \\
 S_2 \leq S_{10} & S_6 \leq S_9 & S_8 \leq \frac{27}{17}S_{10} & S_{10} \leq S_9 \\
 & S_6 \leq S_{10} & &
 \end{array}$$

where $k_4 \approx 1.017624086$ is the smallest positive root of $25947x^7 + 653697x^6 - 49857885x^5 + 128952193x^4 - 112076935x^3 + 32426283x^2 - 3014327x + 2963603$.

We did not check for the conditions when equality occurs.

Corollary 8. *For all triangles,*

$$27r^2 \leq S_4 \leq S_6 \leq S_1 \leq S_2 \leq S_{10} \leq S_9 \leq S_8 \leq 12R^2.$$

Related results can be found in [7].

5. ACUTE TRIANGLES

Algorithm K in [8] allows us to search for inequalities that are true for all acute triangles. We get the following results. We did not check for the conditions when equality occurs.

Theorem 9. *The inequalities given by Theorem 1 are best possible when the triangles are restricted to acute triangles. In addition, the following inequality is true for all acute triangles.*

$$S_3 \leq 9R^2$$

Theorem 10. *The following inequalities are true for all acute triangles.*

$$\begin{aligned} (75 - 28\sqrt{7})s^2 &\leq S_1 \leq s^2 \\ s^2 &\leq S_2 \leq \frac{3}{2}s^2 \\ 1s^2 &\leq S_3 \leq \frac{17}{9}s^2 \\ (15 - 10\sqrt{2})s^2 &\leq S_4 \leq s^2 \\ \frac{49}{9}(3 - 2\sqrt{2})s^2 &\leq S_5 \leq \frac{33}{25}s^2 \\ k_5s^2 &\leq S_6 \leq s^2 \\ \frac{147 - 19\sqrt{57}}{4}s^2 &\leq S_7 \leq s^2 \\ s^2 &\leq S_8 \leq 3s^2 \\ s^2 &\leq S_9 \leq 2s^2 \\ s^2 &\leq S_{10} \leq \frac{17}{9}s^2 \\ \frac{7}{8}s^2 &\leq S_{11} \leq (27 - 18\sqrt{2})s^2 \\ \frac{8043 - 5330\sqrt{2}}{529}s^2 &\leq S_{12} \leq s^2 \end{aligned}$$

where $k_5 \approx 0.8742445769$ is the positive root of $5000x^6 + 32241x^5 + 215799x^4 - 164970x^3 - 239166x^2 + 258633x - 77841$.

Constants in blue are those that differ from the corresponding constant in Theorem 2.

Theorem 11. *The following inequalities are true for all acute triangles.*

$$\begin{aligned} \frac{27}{2}rR &\leq S_1 \leq \frac{3}{2}\sqrt{3}Rs \\ \frac{27}{2}rR &\leq S_2 \leq 3Rs \\ \frac{27}{2}rR &\leq S_3 \leq \frac{34}{9}Rs \\ (5 + 5\sqrt{2})rR &\leq S_4 \leq \frac{3}{2}\sqrt{3}Rs \\ \frac{49 + 49\sqrt{2}}{9}rR &\leq S_5 \leq \frac{66}{25}Rs \\ \frac{49 + 49\sqrt{2}}{9}rR &\leq S_6 \leq \frac{3}{2}\sqrt{3}Rs \\ (3 + 7\sqrt{2})rR &\leq S_7 \leq \frac{3}{2}\sqrt{3}Rs \\ \frac{27}{2}rR &\leq S_8 \leq 6Rs \\ \frac{27}{2}rR &\leq S_9 \leq \frac{34}{9}Rs \\ \frac{27}{2}rR &\leq S_{10} \leq \frac{34}{9}Rs \\ k_2rR &\leq S_{11} \leq (9\sqrt{2} - 9)Rs \\ \frac{3001 + 2905\sqrt{2}}{529}rR &\leq S_{12} \leq \frac{3}{2}\sqrt{3}Rs \end{aligned}$$

where $k_2 \approx 14.12657721$ is the positive root of $2x^3 - 5x^2 - 256x - 1024$.

Constants in blue are those that differ from the corresponding constant in Theorem 3.

Theorem 12. *The inequalities given by Theorem 4 are best possible when the triangles are restricted to acute triangles. In addition, the following inequality is true for all acute triangles.*

$$5K \leq S_4$$

Theorem 13. *The following inequalities are true for all acute triangles.*

$$\begin{aligned} \frac{1}{2}(a^2 + b^2 + c^2) &\leq S_1 \leq \frac{3}{4}(a^2 + b^2 + c^2) \\ S_2 &= \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{1}{2}(a^2 + b^2 + c^2) &\leq S_4 \leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{1}{2}(a^2 + b^2 + c^2) &\leq S_6 \leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{1}{2}(a^2 + b^2 + c^2) &\leq S_7 \leq \frac{3}{4}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_8 \leq \frac{3}{2}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_9 \leq \frac{17}{18}(a^2 + b^2 + c^2) \\ \frac{3}{4}(a^2 + b^2 + c^2) &\leq S_{10} \leq \frac{17}{18}(a^2 + b^2 + c^2) \end{aligned}$$

Constants in blue are those that differ from the corresponding constant in Theorem 5.

Theorem 14. *The following inequalities are true for all acute triangles.*

$$\begin{aligned} \frac{6\sqrt{2}-7}{2}(ab+bc+ca) &\leq S_1 \leq ab+bc+ca \\ \frac{3}{4}(ab+bc+ca) &\leq S_2 \leq \frac{3}{2}(ab+bc+ca) \\ \frac{10\sqrt{2}-5}{14}(ab+bc+ca) &\leq S_4 \leq ab+bc+ca \\ k_6 &\leq S_6 \leq ab+bc+ca \\ \frac{26\sqrt{2}-27}{14}(ab+bc+ca) &\leq S_7 \leq ab+bc+ca \\ \frac{3}{4}(ab+bc+ca) &\leq S_8 \leq 3(ab+bc+ca) \\ \frac{3}{4}(ab+bc+ca) &\leq S_9 \leq \frac{17}{9}(ab+bc+ca) \\ \frac{3}{4}(ab+bc+ca) &\leq S_{10} \leq \frac{17}{9}(ab+bc+ca) \end{aligned}$$

where $k_6 \approx 0.7067084379$ is the positive root of $512000x^7 + 831488x^6 + 519424x^5 - 82176x^4 + 1093104x^3 - 2084400x^2 + 946647x - 233523$.

Constants in blue are those that differ from the corresponding constant in Theorem 6.

REFERENCES

- [1] Oene Bottema, R. Ž. Djordjević, R. R. Janic, Dragoslav S. Mitrinović, and P. M. Vasic, *Geometric Inequalities*, Wolters-Noordhoff Publishing, Groningen, The Netherlands, 1969.
- [2] Sava Grozdev and Deko Dekov, *Barycentric Coordinates: Formula Sheet*, International Journal of Computer Discovered Mathematics, **1**(2016)75–82.
<http://www.journal-1.eu/2016-2/Grozdev-Dekov-Barycentric-Coordinates-pp.75-82.pdf>
- [3] Clark Kimberling, *Central Points and Central Lines in the Plane of a Triangle*, Mathematics Magazine, **67**(1994)163–187.
<https://www.jstor.org/stable/2690608>
- [4] Clark Kimberling, *Encyclopedia of Triangle Centers*.
<http://faculty.evansville.edu/ck6/encyclopedia/ETC.html>
- [5] Martin Lukarevski and Dan Ștefan Marinescu, *An inequality for cevians and applications*, Elemente der Mathematik, **75**(2020)166–171.
<http://doi.org/10.4171/EM/418>
- [6] Dragoslav S. Mitrinović, J. Pečarić, and Vladimir Volenec, *Recent advances in geometric inequalities*. Kluwer Academic Publishers, Dordrecht, The Netherlands, 1989.
https://books.google.com/books?id=_OvGErB0QK4C
- [7] Stanley Rabinowitz. *Inequalities Involving Gergonne and Nagel Cevians*, International Journal of Computer Discovered Mathematics, **6**(2021)78–83.
<http://www.journal-1.eu/2021/Stanley%20Rabinowitz.%20Inequalities%20Involving%20Gergonne%20and%20Nagel%20Cevians,%20pp.%2078-83..pdf>
- [8] Stanley Rabinowitz. *A Computer Algorithm for Proving Symmetric Homogeneous Triangle Inequalities*, International Journal of Computer Discovered Mathematics, **7**(2022)30–62.
<http://www.journal-1.eu/2022/3.%20Stanley%20Rabinowitz.%20A%20Computer%20Algorithm%20for%20Proving%20Symmetric%20Homogeneous%20Triangle%20Inequalities,%20pp.%2030-62..pdf>